

Appendix B

B.1. Freezing of Gait Questionnaire (FOGQ)

B.1.1. During your worst state—Do you walk:

- 0 Normally
- 1 Almost normally—somewhat slow
- 2 Slow but fully independent
- 3 Need assistance or walking aid
- 4 Unable to walk

B.1.2. Are your gait difficulties affecting your daily activities and independence?

- 0 Not at all
- 1 Mildly
- 2 Moderately
- 3 Severely
- 4 Unable to walk

B.1.3. Do you feel that your feet get glued to the floor while walking, making a turn or when trying to initiate walking (freezing)?

- 0 Never
- 1 Very rarely—about once a month
- 2 Rarely—about once a week
- 3 Often—about once a day
- 4 Always—whenever walking

B.1.4. How long is your longest freezing episode?

- 0 Never happened
- 1 1–2 s
- 2 3–10 s
- 3 11–30 s
- 4 Unable to walk for more than 30 s

B. 1.5. How long is your typical start hesitation episode (freezing when initiating the first step)?

- 0 None
- 1 Takes longer than 1 s to start walking
- 2 Takes longer than 3 s to start walking

3 Takes longer than 10 s to start walking

4 Takes longer than 30 s to start walking

B. 1.6. How long is your typical turning hesitation: (freezing when turning)

- 0 None
- 1 Resume turning in 1–2 s
- 2 Resume turning in 3–10 s
- 3 Resume turning in 11–30 s
- 4 Unable to resume turning for more than 30 s

References

- [1] Giladi N, McMahon D, Przedborski S, Flaster E, Guillory S, Kostic V, Fahn S. Motor blocks in Parkinson's disease. *Neurology* 1992;42:333–9.
- [2] Giladi N, Kao R, Fahn S. Freezing phenomenon in patients with parkinsonism. *Mov Disord* 1992;12:302–5.
- [3] Fahn S. The freezing phenomenon in parkinsonism. In: Fahn S, Hallet M, Lüder HO, Marsden CD, editors. *Advances in neurology*, vol. 65. Philadelphia, PA: Lippincott-Raven Publishers, 1995. p. 53–63.
- [4] Nieuwboer A, De Weerd W, Dom R, Lesaffre EA. A frequency and correlation analysis of motor deficits in Parkinson patients. *Disabil Rehabil* 1998;20:142–50.
- [5] Narabayashi H, Kondo T, Nagatsu T, Hayashi A, Suzuki T. DL-threo-3,4-dihydroxyphenylserine for freezing symptom in parkinsonism. In: Hassler RG, Christ JE, editors. *Advances in neurology*, vol. 40. New York: Raven Press, 1984. p. 497–552.
- [6] Achiron A, Ziv I, Goren M, Goldberg H, Zoldan Y, Sroka H, Melamed E. Primary progressive freezing gait. *Mov Disord* 1993;8:293–7.
- [7] Fahn S, Marsden CD, Calne DB, Goldstein M, editors. *Recent developments in Parkinson's disease*, vol. 2. Florham Park, NJ: Macmillan Health Care Information, 1987. p. 153–63 (also p. 293–304).
- [8] Huges AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease—a clinico pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* 1992;55:181–4.
- [9] Hoehn MM, Yahr MD. Parkinsonism: onset, progression and mortality. *Neurology* 1967;17:427–42.
- [10] Burleigh-Jacobs X, Horak FB, Nutt JG, Obeso JA. Step initiation in Parkinson's disease: influence of levodopa and external sensory triggers. *Mov Disord* 1997;12:206–15.
- [11] Stern GM, Lander CM, Lees AJ. Akinetic freezing and trick movements in Parkinson's disease. *J Neural Transm* 1980;16(Suppl.):137–41.